

ANT1A

NOVAptamer to Streptomycin


Target Information

Streptomycin

Streptomycin is an aminoglycoside antibiotic used to treat bacterial infections, particularly those caused by Gram-negative bacteria. It works by binding to the bacterial 30S ribosomal subunit, disrupting protein synthesis and causing bacterial cell death. It is primarily used as part of the multi-drug treatment of pulmonary tuberculosis.

Molecular formula: $C_{21}H_{39}N_7O_{12}$

Molecular weight: 581.5 g/mol



NOVAptamer ANT1A

Chemistry: DNA

Size: 79 nt

Molecular weight: 24546 g/mol

Molar extinction coefficient : $764600 \text{ Lmol}^{-1}\text{cm}^{-1}$

Binding buffer: 20 mM HEPES, 20 mM CH_3COONa , 140 mM CH_3COOK , 3 mM $(\text{CH}_3\text{COO})_2\text{Mg}$, pH 7.4

Folding an aptamer into its tertiary structure is essential for optimal target binding. To achieve this, resuspend the aptamer in assay buffer, heat to 95°C (~2 minutes), then allow to cool to room temperature (~5 minutes) before use.

Affinity Determination

Affinity Determination Method: Fluorescence

Apparent K_D in the binding buffer: 31 μM

Specificity:

- cross reacts with: dihydrostreptomycin.




Figure 1. Dose-response fluorescence recovery in a molecular beacon assay upon Streptomycin addition.

Key advantages offered by aptamers over other affinity reagents, notably antibodies

✓	High affinity and selectivity
✓	Thermostable, long shelf life
✓	Animal- and cell-free selection
✓	Chemical synthesis
✓	Batch to batch reproducibility

Custom synthesis

- **Available at different scales** – upon request, up to 100 nanomoles
- **Various purification modes** – adapted to specific experimental requests
- **Extensive conjugation options for diverse applications:**
 - Grafting: NH₂, SH, biotin, etc.
 - Sensing: fluorescent dyes, nanoparticles, redox groups
 - Cross-linking: click chemistry reagents
- **Molecular beacons** - possible hybridization with a complementary oligonucleotide to form a bimolecular beacon, enabling quantitative detection

Applications (For Research Use Only)

- Biosensing
- Fluorescence-based detection in water or dairy products

Contact information

For more information or inquiries, please contact:

NOVAPTECH
2 avenue Favard, 33170 Gradignan, France
<https://novaptech.com> ♦ contact@novaptech.com ♦ +33 (0) 5 47 74 26 85